

Physics 101

Fall Semester

Second Midterm Exam

Saturday, November 29, 2025

8:00 – 9:30 AM

Student's Name: Serial Number:

Student's Number:Section:

Choose your Instructor's Name:

Instructors: Drs. Al Dosari, Al Jassar, Al Qattan, Al Smadi, Askar, Demir, Salameh,
Zaman

For Instructors use only

Grades:

| # | SP1 | SP2 | SP3 | SP4 | SP5 | LP1 | LP2 | Q1 | Q2 | Q3 | Q4 | Total |
|-----|-----|-----|-----|-----|-----|-----|-----|----|----|----|----|-------|
| | 2 | 2 | 2 | 2 | 2 | 3 | 3 | 1 | 1 | 1 | 1 | 20 |
| Pts | | | | | | | | | | | | |


Important:

1. Answer all questions and problems (No solution = no points).
2. Full mark = 20 points as arranged in the above table.
3. **Give your final answer in the correct units.**
4. Assume $g = 10 \text{ m/s}^2$.
5. Mobiles are **strictly prohibited** during the exam.
6. Programmable calculators, which can store equations, are not allowed.
7. Please box your answers.
8. **Cheating incidents will be processed according to the university rules.**


GOOD LUCK

Part I: Short Problems (2 points each)

SP1. A block of mass $m = 22 \text{ kg}$ is pulled by a constant force \vec{F} up a **frictionless** incline, as shown. If the block moves up the incline at **constant speed**, find the magnitude of the force $|\vec{F}|$.




SP2. Two blocks ($m_1 = 7 \text{ kg}$, $m_2 = 3 \text{ kg}$) are in contact on a horizontal, **frictionless** surface, as shown. A horizontal force $\vec{F} = 30 \text{ N}$ is applied to block 1. Find the magnitude of the force that block 1 exerts on block 2.




SP3. A constant net force $\vec{F} = (3\hat{i} + 4\hat{j}) \text{ N}$ acts on an object that starts moving from **the origin** at $t = 0$ and reaches a position $\vec{r} = (12\hat{i} + 16\hat{j}) \text{ m}$ at $t = 4 \text{ s}$. Find the average power delivered by \vec{F} .

SP4. A box of mass 5 kg , which compresses a spring ($k = 900\text{ N/m}$) by 0.2 m , is released from **rest** such that the spring pushes the box along a **rough** horizontal surface ($\mu_k = 0.45$), as shown. **Find the maximum distance S that the box will travel.** (Note: The box is not attached to the spring).




SP5. A 0.5 kg ball is connected to a light string and rotates in a **vertical circle** of radius 3 m , as shown. If the **string breaks** when the tension reaches its maximum value of 33 N , **what is the maximum speed the ball can have while continuing to rotate?**




Part II: Long Problems (3 points each)

LP1. Box A of mass 10 kg rests on a rough horizontal surface ($\mu_s = 0.4$, $\mu_k = 0.37$) is connected to box B of mass 3 kg by a light string that passes over frictionless light pulley, as shown.

a) Find the static friction force acting on box A .




b) A pulling force $|\vec{F}|$ is applied to box B . Find the magnitude of the minimum pulling force $|\vec{F}|$ required to make the boxes start moving.



c) After the boxes start moving, find the magnitude of the pulling force $|\vec{F}|$ required to make the boxes move at constant speed.

LP2. A small block of mass $m = 0.5 \text{ kg}$ passes through **point A** with a speed of 5 m/s . Its path is frictionless until it reaches the section of length $L = 12 \text{ m}$, where the coefficient of kinetic friction is μ_k . The indicated heights are $h_1 = 4 \text{ m}$ and $h_2 = 2 \text{ m}$.




a) Find the change in the gravitational potential energy as the block moves from A to B.

b) Find the speed of the block at point C.


c) If the block completely stops at point D, find μ_k .

Part III: Questions (Choose the correct answer, one point each)

Q1. You pull horizontally on block B , as shown, causing **both blocks to accelerate together as a unit**. Which of the following correctly represents the free-body diagram of block B if the table is frictionless?




*



*



*



*


Q2. A 30 kg boy stands on a scale while riding in the elevator. If the scale reads 330 N, then the elevator is moving:

- * downward with increasing speed.
- * downward with decreasing speed.
- * upward with decreasing speed.
- * upward with constant speed.



Q3. The work done by a radial force (\vec{F}_{rad}) on an object moving in a circle from point A to point B with increasing speed is:

- * $W = 0$
- * $W = \Delta K$
- * $W = |\vec{F}_{tan}| |\vec{S}|$
- * $W = |\vec{F}_{rad}| |\vec{S}|$



Q4. A particle is under the influence of a net force along the x -axis whose graph is shown. Which of the following statements is true about its initial speed (v_i) at x_i and its final speed (v_f) at x_f ?

- * $v_i > v_f$
- * $v_i < v_f$
- * $v_i = v_f \neq 0$
- * $v_i = v_f = 0$

